



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 37/00, A61K 31/21, C07C 69/34, 69/74	A1	(11) International Publication Number: WO 96/39033 (43) International Publication Date: 12 December 1996 (12.12.96)
(21) International Application Number: PCT/US96/09485 (22) International Filing Date: 5 June 1996 (05.06.96) (30) Priority Data: 08/463,071 5 June 1995 (05.06.95) US (71) Applicants: THE UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL [US/US]; Office of Technology Development, Campus Box #4105, 308 Bynum Hall, Chapel Hill, NC 27599-4105 (US). BIOTECH RESEARCH LABORATORIES [US/US]; 3 Taft Court, Rockville, MD 20850 (US). (72) Inventors: LEE, Kuo-Hsiung; 1426 Gray Bluff Trail, Chapel Hill, NC 27514 (US). KASHIWADA, Yoshiki; 5-14-31, Kamishin'ei-cho, Niigata 950-21 (JP). HASHIMOTO, Fumio; 4-1-11-43, Higashi-machi, Kumamoto 862 (JP). COSENTINO, Louis, Mark; 6648 Green Ash Drive, Springfield, VA 22152 (US). MANAK, Mark; 200 Stanley Place, Laurel, MD 20707 (US). (74) Agents: GOLDSTEIN, Jorge, A. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: BETULINIC ACID DERIVATIVES AND USES THEREFOR**(57) Abstract**

Some betulinic acid and dihydrobetulinic acid acyl derivatives according to the present invention have been found to have potent anti-HIV activity. Introducing a C₂-C₂₀ substituted or unsubstituted acyl group at the C₃-hydroxy group of betulinic acid and dihydrobetulinic acid produces the corresponding 3-O-acyl derivatives. The compounds of the present invention have formulae (I) and (II), where R may be a mono- or dicarboxyacyl group, substituted or unsubstituted, of from about 2 to about 20 carbon atoms, and R' may be hydrogen or a C₂-C₁₀ substituted and unsubstituted alkyl or aryl group.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

BETULINIC ACID DERIVATIVES AND USES THEREFOR

Field of the Invention

5 The present invention relates to betulinic acid and dihydrobetulinic acid and derivatives thereof and their use as pharmaceuticals.

Background of the Invention

Retroviruses

10 Retroviruses are small, single-stranded positive-sense RNA viruses. A retroviral particle comprises two identical single-stranded positive sense RNA molecules. Their genome contains, among other things, the sequence of the RNA-dependent DNA polymerase, also known as reverse transcriptase. Many molecules of reverse transcriptase are found in close association with the genomic RNA in the mature viral particles. Upon entering a cell, this reverse transcriptase produces a double-stranded DNA copy of the viral genome, which is then inserted
15 into the chromatin of a host cell. Once inserted, the viral sequence is called a provirus. Retroviral integration is directly dependent upon viral proteins. Linear viral DNA termini (the LTRs) are the immediate precursors to the integrated proviral DNA. There is a characteristic duplication of short stretches of the host's DNA at the site of integration.

20 Progeny viral genomes and mRNAs are transcribed from the inserted proviral sequence by host cell RNA polymerase in response to transcriptional, regulatory signals in the terminal regions of the proviral sequence, the long terminal repeats, or LTRs. The host cell's protein production machinery is used to produce viral proteins, many of which are inactive until processed by virally
25 encoded proteases. Typically, progeny viral particles bud from the cell surface in a non-lytic manner. Retroviral infection does not necessarily interfere with the normal life cycle of an infected cell or organism. However, neither is it always

-2-

benign with respect to the host organism. While most classes of DNA viruses can be implicated in tumorigenesis, retroviruses are the only taxonomic group of RNA viruses that are oncogenic. Various retroviruses, such as the Human Immunodeficiency Virus (HIV), which is the etiological agent responsible for acquired immune deficiency syndrome (AIDS) in humans, are also responsible for several very unusual diseases of the immune system of higher animals.

HIV Infection and AIDS

Human Immunodeficiency Virus (HIV) is a member of the lentiviruses, a subfamily of retroviruses. Many retroviruses are well-known carcinogens. HIV *per se* is not known to cause cancer in humans or other animals, but it does present a formidable challenge to the host. The viral genome contains many regulatory elements which allow the virus to control its rate of replication in both resting and dividing cells. Most importantly, HIV infects and invades cells of the immune system; it breaks down the body's immune system and renders the patient susceptible to opportunistic infections and neoplasms. The immune defect appears to be progressive and irreversible, with a high mortality rate that approaches 100% over several years.

HIV-1 is trophic and cytopathic for T4 lymphocytes, cells of the immune system which express the cell surface differentiation antigen CD4, also known as OKT4, T4 and leu3. The viral tropism is due to the interactions between the viral envelope glycoprotein, gp120, and the cell-surface CD4 molecules (Dalglish *et al.*, *Nature* 312:763-767 (1984)). These interactions, not only mediate the infection of susceptible cells by HIV, but are also responsible for the virus-induced fusion of infected and uninfected T cells. This cell fusion results in the formation of giant multinucleated syncytia, cell death, and progressive depletion of CD4 cells in AIDS patients. These events result in HIV-induced immunosuppression and its subsequent sequelae, opportunistic infections and neoplasms.

-3-

In addition to CD4+ T cells, the host range of HIV includes cells of the mononuclear phagocytic lineage (Dalglish *et al.*, *supra*), including blood monocytes, tissue macrophages, Langerhans cells of the skin and dendritic reticulum cells within lymph nodes. HIV is also neurotropic, capable of infecting monocytes and macrophages in the central nervous system causing severe neurologic damage. Macrophage/monocytes are a major reservoir of HIV. They can interact and fuse with CD4-bearing T cells, causing T cell depletion and thus contributing to the pathogenesis of AIDS.

Anti-HIV Drugs

Intensive efforts are currently under way to develop therapies to prevent or intervene in the development of clinical symptoms in HIV-infected individuals. For the most part, efforts have been focused on the use of nucleoside analogue drugs such as AZT (azidothymidine), and on other dideoxynucleoside derivatives such as ddA, ddT, ddI, and ddC. These drugs inhibit the viral enzyme reverse transcriptase, thereby inhibiting *de novo* infection of cells. However, once viral infection has been established within a cell, viral replication utilizes host cell enzymes. Thus, drugs which inhibit only reverse transcriptase tend to have limited effects. While the spread of free virus within the organism can be blocked, the mechanisms of syncytia formation and pathogenesis through direct intercellular spread remain. Accordingly, there is a need to provide new anti-HIV drugs which are not limited to inhibiting reverse transcription as their mechanism of action.

Various efforts are currently underway to develop therapeutic agents to arrest the replication of the human immunodeficiency virus (HIV). Although several nucleoside HIV-1 reverse transcriptase (RT) inhibitors, including AZT, ddI, ddC and D4T have been approved by the FDA and are in use clinically, all of these compounds have adverse side effects, and the HIV mutates so quickly that the virus has developed resistance to these drugs. Therefore, compounds

which possess potent anti-HIV activity with different modes of action are urgently needed to add to existing anti-HIV therapies. Currently, development of new anti-HIV agents is focused on discovering different compounds which either have novel structures or new mechanisms of action.

5 Much effort has been expended on plant-derived natural products as new lead compounds for anti-HIV agents, as well as developing modifications of these lead compounds. Suitable structural modification of the initial lead structures may provide derivatives with greatly enhanced activity. As one example, some of the present inventors isolated and identified suksdorfin as an anti-HIV
10 compound from *Lomatium suksdorfii*. Subsequent modification of suksdorfin yielded 3',4'-di-O-(-)-camphanoyl-(+)-*cis*-khellactone (DCK), which compound has demonstrated extremely potent inhibitory activity against HIV replication in H9 lymphocyte cells with an EC₅₀ value of 0.0004 μ M and a therapeutic index (T.I.) value of 136,719.

15 Previously, betulinic acid and platanic acid were isolated as anti-HIV principles from *Syzigium claviflorum*. Betulinic acid (1) and platanic acid (12) exhibited inhibitory activity against HIV-1 replication in H9 lymphocyte cells with EC₅₀ values of 1.4 μ M and 6.5 μ M, respectively, and T.I. values of 9.3 and 14, respectively. Hydrogenation of betulinic acid yielded dihydrobetulinic acid
20 (6), which showed slightly more potent anti- HIV activity with an EC₅₀ value of 0.9 and a T.I. value of 14. Based upon these findings, modification of these lead compounds, betulinic acid and dihydrobetulinic acid, has been carried out and has led to the discovery of additional potent anti-HIV agents.

25 A number of triterpenoids, including betulinic acid, have several known medical applications, including use as an anticancer drug. Anderson *et al.*, in WO 95/04526, discuss derivatives of triterpenoids which have been used in cancer therapy, including their activity against polyamines which are required by cells to grow at an optimal rate. Some of these triterpenoids have been found to interfere with the enzymatic synthesis of polyamines required for optimum cell
30 growth, and thus inhibit the growth of cancer cells, particularly by inhibiting

-5-

ornithine decarboxylase. Betulinic acid has been reported also to possess anti-inflammatory activity, which may be due to its capacity to inhibit enzymes involved in leukotriene biosynthesis, including 5-lipoxygenase.


Chan, in British patent 1,425,601, discloses that carboxylated derivatives of dihydrobetulinic acid can be included in pharmaceutical compositions, but there is no hint as to what use these compounds have.

Summary of the Invention

Betulinic acid and dihydrobetulinic acid acyl derivatives according to the present invention have been found to have potent anti-HIV activity. The C₃-hydroxy, C₁₇-carboxylic acid and C₂₀-exomethylene groups in betulinic acid can be easily modified. It has been found that introducing a C₂ to C₂₀ substituted or unsubstituted acyl group at the C₃-hydroxy group of betulinic acid and dihydrobetulinic acid can be readily effective to produce the corresponding 3-O-acyl derivatives. Thus, betulinic acid and dihydrobetulinic acid were treated with 3,3-dimethylglutaric anhydride or diglycolid anhydride in pyridine in the presence of dimethylaminopyridine to produce the corresponding 3-O-acyl derivatives (4, 5, 9 and 10). In contrast, similar treatment of betulinic acid and dihydrobetulinic acid with dimethylsuccinic anhydride produced a mixture of 3-O- (2',2'-dimethylsuccinyl) and 3-O-(3',3' dimethylsuccinyl)-betulinic acid (2 and 3) and dihydrobetulinic acid (7 and 8), respectively. The mixture was successfully separated by preparative scale HPLC yielding pure samples. The structures of these isomers were assigned by long-range ¹H-¹³C COSY examinations.


-6-

The compounds of the present invention have the following formulae:



where R may be a mono- or dicarboxyacyl group, substituted or unsubstituted, of from about 2 to about 20 carbon atoms and R' may be H or a C₂-C₁₀ substituted or unsubstituted alkyl or aryl group, such as acetyl, benzyl, etc.

Of particular interest are compounds of the formula:



Compound

R Group


Compound

1

H


6

2



7




3




8

10

-7-



11



12 (RPR103611)

Anti-HIV assays indicated that 3-O-(3',3'-dimethylsuccinyl)-betulinic acid (3) and -dihydrobetulinic acid (8) both demonstrated extremely potent anti-HIV activity in acutely infected H9 lymphocytes with EC_{50} values of less than 1.7×10^{-5} μ M, respectively. These compounds exhibited remarkable T.I. values of more than 970,000 and more than 400,000. In contrast, compounds 2 and 7, the 2',2'-dimethyl isomers, showed anti-HIV activities with EC_{50} values of 2.7 and

0.56 μ M, respectively, and T.I. values of 5.9 and 13.8, respectively, which were significantly lower than those of 3 and 8. Compounds 4, 5, 9, and 10 also exhibited potent anti-HIV activities with EC_{50} ranging from 0.01 to 2.3×10^{-3} μ M, and T.I. values from 1017 to 2344.

5 The C_3 acyl groups of the more active compounds have dimethyl groups or oxygen at the C3 position. Since the lone pairs of the oxygen in the diglycoyl group might correspond to the dimethyl groups at C3, in the dimethylsuccinyl or dimethylglutaryl groups, these three acyl groups are structurally similar to one another. This observation suggests that this type of acyl group might be
10 important to the enhanced anti-HIV activity.

 The inhibitory activities of 8 and 9 against HIV-1 replication in PHA-stimulated peripheral blood mononuclear cells (PBMCs) were also evaluated. These compounds also displayed potent inhibitory effects in this assay with EC_{50} values of 0.00299 and 0.0292 μ M, respectively, although the T.I. values for these
15 compounds in PBMCs were lower than those in H9 cells.

 As a mechanism of action study, the inhibitory activity of compounds 1-6 and 8-10 against HIV-1 reverse transcriptase was investigated. The tested compounds did not inhibit HIV reverse transcriptase activity at a concentration of 100 μ g/ μ L. Recently, other betulinic acid derivatives, e.g., RPR103600, have
20 been reported as anti-HIV agents. They were shown to inhibit HIV-induced membrane fusion. Therefore, compounds 1-6 and 8-10 were also evaluated for inhibitory activity against HIV induced membrane fusion. Compounds 2-5 and 8-10 inhibited syncytia formation in a concentration range of 20-40 μ g/mL, suggesting than an inhibitory effect against HIV induced membrane fusion could
25 be involved in their mechanisms of action. However, the anti-HIV activity of compound 3 is 540,000 times greater than that of compound 2, although these compounds inhibited syncytia formation at the same concentration. Compound RPR103611 was reported to inhibit syncytia formation at 3 μ g/mL and for this purpose was approximately ten times more potent than compounds 2-5 or 8-10
30 in this assay. However, the cell line used in assaying the compounds of the

present invention, CEM4, was different from the cell line used in the literature, H9. The anti-HIV activities of compounds 3 and 8 in H9 cells and PMBCs were about 10 to 1000 times greater than those of RPR103611 reported in the literature. This observation indicated that at least one other mechanism of action other than inhibition of syncytia formation could be involved in the anti-HIV activity shown by these compounds.

The invention is also directed to a method for treating a subject infected with HIV-1 by administering at least one of the above-noted betulinic acid derivatives, optionally in combination with any one or more of the known anti-AIDS therapeutics or an immunostimulant.

The treatment methods of the present invention also include administering to a subject infected with HIV-1 a conjugate of a betulinic acid derivative as described above with soluble CD4, CD4 derivatives, antibodies specific for CD4, or HIV-coded glycoproteins such as gp120 and gp41, or antibodies thereto.

Other features, advantages, embodiments, aspects and objects of the present invention will be clear to those skilled in the areas of relevant art, based upon the description, teaching and guidance presented herein.

The compounds of the present invention have been unexpectedly discovered to have anti-retroviral activity, thus providing suitable compounds and compositions for treating retroviral infections, optionally with additional pharmaceutically active ingredients, such as anti-retroviral, anti-HIV, and/or immuno-stimulating compounds or antiviral antibodies or fragments thereof.

By the term "anti-retroviral activity" or "anti-HIV activity" is intended the ability to inhibit at least one of:

- (1) retroviral attachment to cells;
- (2) viral entry into cells;
- (3) cellular metabolism which permits viral replication;
- (4) inhibition of intercellular spread of the virus;
- (5) synthesis and/or cellular expression of viral antigens;

-10-

(6) activity of virus-coded enzymes (such as reverse transcriptase and protease); and/or

(7) any known retroviral or HIV pathogenic actions, such as, for example, immunosuppression. Thus, any activity which tends to inhibit any of these mechanisms is "anti-retroviral activity" or "anti-HIV activity."

Detailed Description of the Preferred Embodiments

A betulinic acid or dihydrobetulinic acid derivative of the present invention can be used for treatment of retroviral (e.g., HIV) infection either alone, or in combination with other modes of therapy known in the art. Such modes of therapy can include chemotherapy with drugs, such as, but not limited to, at least one of AZT, ddC, ddA, ddT, ddI, or any other antiretroviral drugs or antibodies in combination with each other, or associated with a biologically based therapeutic, such as, for example, soluble CD4, antibodies to CD4, and conjugates of CD4 or anti-CD4, or as additionally presented herein.

Because the betulinic acid or dihydrobetulinic acid derivatives of the present invention are relatively less or substantially non-toxic to normal cells, their utility is not limited to the treatment of established retroviral infections. For example, a betulinic acid derivative according to the present invention can be used in treating blood products, such as those maintained in blood banks. The nation's blood supply is currently tested for antibodies to HIV. However, the test is still imperfect and samples which yield negative tests can still contain HIV virus. Treating the blood and blood products with the betulinic acid derivatives of the present invention can add an extra margin of safety by killing any retrovirus that may have gone undetected.

Pharmaceutical Compositions

Pharmaceutical compositions of the present invention can comprise at least one of the betulinic acid or dihydrobetulinic derivatives. Pharmaceutical compositions according to the present invention can also further comprise other anti-viral agents such as, but not limited to, AZT, ddI, 2'- β -fluoro-ddI, ddA, ddG, ddC, 2'- β -fluoro-ddC, d4T, AzddU, phosphonylmethoxyethyl-adenine, or soluble CD4, or immunomodulators, e.g., as presented below. For a review of therapeutic agents in HIV infections, see, e.g., Mitsuya *et al.*, *FASEB J.* 5:2369-2381 (1991), which reference is hereby incorporated by reference.

Additional suitable antiviral agents for optimal use with a betulinic acid derivative of the present invention can include, but are not limited to, AL-721 (lipid mixture) manufactured by Ethigen Corporation and Matrix Research Laboratories; Amphotericin B methyl ester; Ampligen (mismatched RNA) developed by DuPont/HEM Research; anti-AIDS antibody (Nisshon Food) 1 AS-101 (heavy metal based immunostimulant); AZT (azidothymidine/Retrovir/Zidovudine), manufactured by Burroughs Wellcome; Betaseron (β -interferon) manufactured by Triton Biosciences (Shell Oil); butylated hydroxytoluene; Carrosyn (polymannoacetate) Castanospermine; Contracan (stearic acid derivative); Creme Pharmatex (containing benzalkonium chloride) manufactured by Pharmalec; CS-87 (5-unsubstituted derivative of Zidovudine), Cytovene (ganciclovir) manufactured by Syntex Corporation; ddC (dideoxycytidine) manufactured by Hoffmann-LaRoche and other nucleoside analogues; dextran sulfate; D-penicillamine (3-mercapto-D-valine) manufactured by Carter-Wallace and Degussa Pharmaceutical; Foscarnet (trisodium phosphonoformate) manufactured by Astra AB, fusidic acid manufactured by Leo Lovens, glycyrrhizin (a constituent of licorice root); HPA-23 (ammonium-21-tungsto-9-antimonate) manufactured by Rhone-Poulenc Santé; human immune virus antiviral developed by Porton Products International; Ornidyl (eflornithine) manufactured by Merrell-Dow; nonoxinol; pentamidine isethionate (PENTAM-

300) manufactured by Lypho Med; Peptide T (octapeptide sequence) manufactured by Peninsula Laboratories; Phenytoin (Warner-Lambert); Ribavirin; Rifabutin (ansamycin) manufactured by Adria Laboratories; rsT4 (recombinant soluble T4) manufactured by Biogen, Genentech and Smith, Kline Beecham; Trimetrexate manufactured by Warner-Lambert Company; SK-818 (germanium-derived antiviral) manufactured by Sanwa Kagaku; suramin and analogues thereof manufactured by Miles Pharmaceuticals; UA001 manufactured by Ueno Fine Chemicals Industry; Wellferon (α -interferon) manufactured by Burroughs Wellcome; Zovirex (acyclovir, AZT) manufactured by Burroughs Wellcome.

Pharmaceutical compositions of the present invention can also further comprise immunomodulators. Suitable immunomodulators for optional use with a betulinic acid derivative of the present invention in accordance with the present invention can include, but are not limited to: ABPP (Bropiramine); Ampligen (mismatched RNA) DuPont/HEM Research; anti-human interferon- α -antibody (Advance Biotherapy and Concepts); anti-AIDS antibody (Nisshon Food); AS-101 (heavy metal based immunostimulate; ascorbic acid and derivatives thereof; interferon- β ; Carrosyn (polymannoacetate); Ciamexon (Boehringer-Mannheim); cyclosporin; cimetidine; CL-246,738 (American Cyanamide); colony stimulating factors, including GM-CSF (Sandoz, Genetics Institute); dinitrochlorobenzene; interferon- α ; inteferon-gamma; glucan; hyperimmune gamma-globulin (Bayer); IMREG-1 (leukocyte dialyzate) and IMREG-2 (IMREG Corp.); immuthiol (sodium diethylthiocarbamate) (Institut Merieux); interleukin-1 or interleukin-1 (Cetus Corporation; Hoffmann-LaRoche; Immunex); isoprinosine (inosine pranobex); Krestin(Sankyo); LC-9018 (Yakult); lentinan (Ajinomoto/Yamanouchi); LF-1695 (Fournier); methionine-enkephalin (TNI Pharmaceuticals; Sigma Chemicals); Minophagen C; muramyl tripeptide, MTP-PE (Ciba-Geigy); naltrexone ("Trexan" DuPont); Neutropin, RNA immunomodulator (Nippon Shingaku); shosaikoto and ginseng; thymic humoral factor; TP-05 (Thymopentin, Ortho Pharmaceuticals); Thymosin factor 5 and

Thymosin 1; Thymostimulin; TNF (Tumor necrosis factor) manufactured by Genentech; and vitamin B preparations.

5 The preferred animal subject of the present invention is a mammal. By the term "mammal" is meant an individual belonging to the class Mammalia. The invention is particularly useful in the treatment of human patients.

The term "treating" means the administering to subjects a betulinic acid or dihydrobetulinic acid derivative for purposes which can include prevention, amelioration, or cure of a retroviral-related pathology.

10 Medicaments are considered to be provided "in combination" with one another if they are provided to the patient concurrently or if the time between the administration of each medicament is such as to permit an overlap of biological activity.

In one preferred embodiment, at least one betulinic acid or dihydrobetulinic acid derivative comprises a single pharmaceutical composition.

15 Pharmaceutical compositions for administration according to the present invention can comprise at least one betulinic acid or dihydrobetulinic acid derivative according to the present invention in a pharmaceutically acceptable form optionally combined with a pharmaceutically acceptable carrier. These compositions can be administered by any means that achieve their intended purposes. Amounts and regimens for the administration of a betulinic acid derivative according to the present invention can be determined readily by those with ordinary skill in the clinical art of treating a retroviral pathology.

20 For example, administration can be by parenteral, such as subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, or buccal routes. Alternatively, or concurrently, administration can be by the oral route. The dosage administered depends upon the age, health and weight of the recipient, type of previous or concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

25 Compositions within the scope of this invention include all compositions comprising at least one betulinic acid or dihydrobetulinic acid derivative

30

according to the present invention in an amount effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the art. Typical dosages comprise about 0.1 to about 100 mg/kg body weight. The preferred dosages comprise about 1 to about 100 mg/kg body weight of the active ingredient. The most preferred dosages comprise about 10 to about 100 mg/kg body weight.

Therapeutic administration can also include prior, concurrent, subsequent or adjunctive administration of at least one additional betulinic acid or dihydrobetulinic acid derivative according to the present invention or other therapeutic agent, such as an anti-viral or immune stimulating agent. In such an approach, the dosage of the second drug can preferably be the same as or different from the dosage of the first therapeutic agent. Preferably, the drugs are administered on alternate days in the recommended amounts of each drug.

Administration of a compound of the present invention can also optionally include previous, concurrent, subsequent or adjunctive therapy using immune system boosters or immunomodulators. In addition to the pharmacologically active compounds, a pharmaceutical composition of the present invention can also contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Preferably, the preparations, particularly those preparations which can be administered orally and which can be used for the preferred type of administration, such as tablets, dragees, and capsules, and also preparations which can be administered rectally, such as suppositories, as well as suitable solutions for administration by injection or orally, contain from about 0.01 to 99 percent, preferably from about 20 to 75 percent of active compound(s), together with the excipient.

Pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus,

pharmaceutical preparations for oral use can be obtained by combining the active compounds with solid excipients, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

5 Suitable excipients are, e.g., fillers such as saccharide, for example, lactose or sucrose, mannitol or sorbitol; cellulose preparations and/or calcium phosphates, such as tricalcium phosphate or calcium hydrogen phosphate; as well as binders such as starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose,
10 hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents can be added such as the above-mentioned starches and also carboxymethyl starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example,
15 silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions can be used, which can optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide,
20 lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropylmethyl cellulose phthalate are used. Dyestuffs or pigments can be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations
25 of active compound doses.

 Other pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules which can be mixed with fillers such
30 as lactose, binders such as starches, and/or lubricants such as talc or magnesium

stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils or liquid paraffin. In addition, stabilizers can be added.

5 Possible pharmaceutical preparations which can be used rectally include, for example, suppositories which consist of a combination of the active compounds with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules which consist of a combination of the active compounds with a base. Possible base materials include, for example,
10 liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons.

Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts. In addition, suspensions of the active compounds as appropriate oily injection suspensions can be administered. Suitable lipophilic solvents or
15 vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides. Aqueous injection suspensions that can contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension can also contain stabilizers.

20 A pharmaceutical formulation for systemic administration according to the invention can be formulated for enteral, parenteral or topical administration. Indeed, all three types of formulation can be used simultaneously to achieve systemic administration of the active ingredient.

Suitable formulations for oral administration include hard or soft gelatin capsules, dragees, pills, tablets, including coated tablets, elixirs, suspensions,
25 syrups or inhalations and controlled release forms thereof.

Solid dosage forms in addition to those formulated for oral administration include rectal suppositories.

The betulinic acid or dihydrobetulinic acid derivatives of the present
30 invention can also be administered in the form of an implant when compounded

-17-

with a biodegradable slow-release carrier. Alternatively, the betulinic acid or dihydrobetulinic acid derivatives of the present invention can be formulated as a transdermal patch for continuous release of the active ingredient.

5 Suitable formulations for topical administration include creams, gels, jellies, mucilages, pastes and ointments. Suitable injectable solutions include intravenous subcutaneous and intramuscular injectable solutions. Alternatively, the betulinic acid or dihydrobetulinic acid derivatives may be administered in the form of an infusion solution or as a nasal inhalation or spray.

10 Having now generally described the invention, the same will be more readily understood through reference to the following:

15 The derivatives of betulinic acid and dihydrobetulinic acid of the present invention were all synthesized by refluxing a solution of betulinic acid (1) or dihydrobetulinic acid (6), dimethylaminopyridine (1 equivalent mol), and an appropriate anhydride (2.5-10 equivalent mol) in anhydrous pyridine (5-10 mL). The reaction mixture was then diluted with ice water and extracted with CHCl_3 . The organic layer was washed with water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was chromatographed using silica gel column or semi-preparative-scale HPLC to yield the product.

Example 1

20 ***3-O-(2',2'-dimethylsuccinyl) betulinic acid (2)***

25 **3-O-(2',2'-dimethylsuccinyl)betulinic acid (2)** was prepared as above using a solution of betulinic acid, dimethylaminopyridine, and dimethylsuccinic anhydride in anhydrous pyridine. The yield was 3.1%. Crystallization from methanol gave colorless needles, melting point 279-280°C; $[\alpha]_D^{19} +36.2^\circ$ [$c=0.35$, CHCl_3 , -MeOH (1:1)]; positive FABMS m/z 585 ($M+H$)⁺; Negative FABMS m/z 583 ($M-H$)⁻; HR-FABMS calculated for $\text{C}_{36}\text{H}_{57}\text{O}_6$ 585.4155, found m/z 585.4156; ^1H NMR (pyridine- d_5): 0.75, 0.93, 1.03 (x2), 1.06 (each 3H, s; 4-

-18-

(CH₃)₂, 8-CH₃, 10-CH₃, 14-CH₃), 1.49 (6H, 2, 2' - CH₃x2), 1.80 (3H, s, 20-CH₃), 2.94 (2H, s, H2-3'), 3.55 (1H, m, H-19), 4.77 (1H, dd, J=5, 11.5 Hz, H-3), 4.79, 4.95 (each 1H, br s, H-30).

Example 2

5 3-O-(3',3'-dimethylsuccinyl)-betulinic acid (3)

Yield 70% (starting with 542 mg of 1); crystallization from MeOH gave colorless needles; mp 274-276°C; [α]_D¹⁹ +23.5° (c = 0.71), CHCl₃-MeOH [1:1]); Positive FABMS m/z 585 (M+H)⁺; Negative FABMS m/z 583 (M-H)⁻; HR-FABMS calcd for C₃₆H₅₇O₆ 585.4155, found m/z 585.4161; ¹H NMR (pyridine-d₅): 0.73, 0.92, 0.97, 1.01, 1.05 (each 3H, s; 4-(CH₃)₂, 8-CH₃, 10-CH₃, 14-CH₃), 1.55 (6H, s, 3' -CH₃ x 2), 1.80 (3H, s, 20-CH₃), 2.89, 2.97 (each 1H, d, J=15.5 Hz, H-2'), 3.53 (1H, m, H-19), 4.76 (1H, dd, J=5.0, 11.5 Hz, H-3), 4.78, 4.95 (each 1H, br s, H-30).

Example 3

15 3-O-(3',3'-dimethylglutaryl)-betulinic acid (4)

Yield 59.50% (starting with 51.3 mg of 2); crystallization from MeOH-H₂O gave colorless needles; mp 214-215°C; [α]_D²⁰ + 9.8° (c=1.2, CHCl₃-MeOH [1:1]); ¹H-NMR (pyridine-d₅): 0.77, 0.91, 0.95, 1.03, 1.07, 1.80 (each 3H, s; 4-(CH₃)₂, 8-CH₃, 10-CH₃, 14-CH₃, 20-CH₃), 4.73 (1H, dd, J=4.5, 11.5 Hz, H-3), 4.78, 4.95 (each 1H, br.s; H-30). Anal. Calcd for C₃₇H₅₈O₆•4H₂O: C 66.24, H 9.92; found C 65.91, H 9.76.

Example 4

3-O-diglycolyl-betulonic acid (5)

Yield 84.7% (starting with 49.6 mg of 1); an amorphous powder; $[\alpha]_D^{20} + 2.1^\circ$ ($c=1.1$, CHCl_3 -MeOH [1:1]); $^1\text{H-NMR}$ (methanol - d_4 - CDCl_3 [1:1]): 0.85, 0.88 (x 2), 0.97, 1.00 (each 3H, s; 4- $(\text{CH}_3)_2$ 8- CH_3 , 10- CH_3 , 14- CH_3), 1.69 (20- CH_3), 4.21, 4.23 (each 2H, s; H_2 -2' and 4'), 4.65-4.75 (1H, br s, H-3), 4.60, 4.73 (each 1H, br s, H-30). Anal. Calcd for $\text{C}_{34}\text{H}_{52}\text{O}_7 \cdot 4\text{H}_2\text{O}$: C 69.12, H 9.21; found C 69.60, H 9.03.

Example 5

3-O-(2',2'-dimethylsuccinyl)-dihydrobetulonic acid (7)

Yield 4.8% (starting with 155.9 mg of 6); crystallization from MeOH- H_2O gave colorless needles; mp 297 - 298°C ; $[\alpha]_D^{17} - 32.2^\circ$ ($c=0.21$, CHCl_3 -MeOH[1:1]); Positive FABMS m/z 587 ($\text{M}+\text{H}^+$); Negative FABMS m/z 585 ($\text{M}-\text{H}$); HR-FABMS calcd for $\text{C}_{36}\text{H}_{59}\text{O}_6$ 587.4311, found m/z 587.4308; $^1\text{H NMR}$ (pyridine- d_5): 0.85, 0.94 (each 3H, d, $J=6.5$ Hz; 20- $(\text{CH}_3)_2$ 0.77, 0.94, 1.03 (x 2), 1.04 (each 3H, s; 4- $(\text{CH}_3)_2$, 8- CH_3 , 10- CH_3 , 14- CH_3), 1.50 (6H, s, 2' - CH_3 x 2), 2.95 (2H, s, H_2 -3'), 4.79 (1H, dd, $J=4.5$, 11.5 Hz; H-3).

Example 6

3-O-(3',3'-dimethylsuccinyl)-dihydrobetulonic acid (8)

Yield 24.5% (starting with 155.9 mg of 6) crystallization from MeOH- H_2O gave colorless needles; mp 291 - 292°C ; $[\alpha]_D^{20} - 13.4^\circ$ ($c=1.1$, CHCl_3 - MeOH [1:1], $^1\text{H NMR}$ (pyridine- d_5): 0.85, 0.94 (each 3H, d, $J=7.0$ Hz; 20- $(\text{CH}_3)_2$), 0.75, 0.93, 0.97, 1.01, 1.03 (each 3H, s; 4- $(\text{CH}_3)_2$, 8- CH_3 , 10- CH_3 ,

-20-

14-CH₃), 1.55 (6H, s; 3' -CH₃ x 2), 2.89, 2.97 (each 1H, d, $J=15.5$ Hz; H-2'), 4.77 (1H, dd, $J=5.0, 11.0$ Hz, H- 3); Anal. Calcd for C₃₆H₅₈O₆•5/2H₂O: C 68.43, H 10.04; found C 68.64, H 9.78.

Example 7

5 3-O-(3',3'-dimethylglutaryl)-dihydrobetulinic acid (9)

Yield 93.3% (starting with 100.5 mg of 6) ; crystallization from needles MeOH-H₂O gave colorless needles; mp 287 - 289°C; $[\alpha]_D^{20}$ - 17.9% ($c=0.5$, CHCl₃-MeOH [1:1]); ¹H-NMR (pyridine-d₅): 0.86, 0.93 (each 3H, d, $J=6.5$ Hz; 20- (CH₃)₂), 0.78, 0.92, 0.96, 1.02, 1.05 (each 3H, s; 4- (CH₃)₂, 8-CH₃, 10-CH₃, 14-CH₃), 1.38, 1.39 (each 3H, s; 3' - CH₃ x 2) , 2.78 (4H, m, H₂-2' and 4'), 4.76 (1H, dd, $J=4.5, 11.5$ Hz; H-3). Anal. Calcd for C₃₇H₆₀O₆: C 73.96, H 10.06; found C 73.83, H 10.10.

Example 8

3-O-diglycolyl-dihydrobetulinic acid (10)

15 Yield 79.2% (starting with 103.5 mg of 6); an off-white amorphous powder; $[\alpha]_D^{20}$ - 9.8° ($c=1.1$, CHCl₃-MeOH[1:1]); ¹H-NMR (pyridine-d) : 0.79, 0.87 (each 3H, d, $J=6.5$ Hz; 20- (CH₃)₂ 0.87, 0.88, 0.91, 0.98, 1.01 (each 3H, s; 4- (CH₃)₂) 8-CH₃, 10-CH₃, 14-CH₃), 4.21, 4.23 (each 2H, s, H₂-2' and 4'), 4. 57 (1H, dd, $J=6. 5, 10.0$ Hz, H-3); Anal. Calcd for C₃₄H₅₄O₇•2H₂O: C 66.85, H 9.57; found C 67.21, H 9.33.

Example 9

Pharmacological Activity

Anti-HIV Assays

5 The T cell line, H9, and the promonocytic cell line, U937, were maintained separately in continuous culture with complete medium (RPMI 1640 with 10% fetal calf serum) at 5% CO₂ and 37°C. The cell lines were used in experiments only when in the logarithmic phase of growth, whereas uninfected peripheral blood mononuclear cells (PBMCs) were first stimulated with PHA (1 µg/mL) for three days. All cell targets were incubated with HIV-1 (IIIB isolate, 10 TCID₅₀ 10⁴ IU/ml, at a multiplicity of infection of 0.01-0.01 IU/cell) for one hour at 37°C and 5% CO₂. The cell lines and PBMCs were washed thoroughly to remove unadsorbed virions and resuspended at 4 x 10⁵ cells/ml in complete medium or complete medium with 10% v/v interleukin 2 (IL-2), respectively. One ml. aliquots were placed into wells of 24-well culture plates containing an equal volume of test compounds (diluted in the appropriate culture medium). The toxicity of each compound was assessed by determining the number of compound-exposed uninfected cells that remained after four days at 37°C and 5% CO₂. A p24 antigen ELISA assay was used to determine the level of virus released in the medium of the HIV-infected cultures. The p24 antigen assay used 20 a HIV-1 anti-p24 specific monoclonal antibody as the capture antibody coated onto 96-well plates. Following a sample incubation period, rabbit serum containing antibodies for HIV-1 p24 was used to tag any p24 "captured: onto the microtiter well surface. Peroxidase conjugated goat anti-rabbit serum was then used to tag HIV-1 p24 specific rabbit antibodies that had complexed with 25 captured p24. The presence of p24 in test samples was then revealed by addition of substrate. The cutoff for the p24 ELISA assay was 12.5 pg/ml. P24 in the culture medium was quantitated against a standard curve containing known

-22-

amounts of p24. The effective (EC_{50}) and inhibitory (IC_{50}) concentrations for anti-HIV activity and cytotoxicity, respectively, were determined.

HIV-1 Reverse Transcriptase Assay

5 HIV-1 reverse transcriptase microassay was adapted from techniques described by Goff *et al.*, *J. Virol.* 38:239-248 (1981) and Willey *et al.*, *J. Virol.* 62:139-147 (1988).

10 Ten microliters of virion associated HIV-1 IIIB reverse transcriptase in 1% Triton X-100 was mixed with 50 microliters of a reaction cocktail containing 50 mM Tris-HCl (pH 7.8), 75 mM KCl, 2 mM dithiothreitol, 5 mM MgCl, 5 mg/ml poly (A) [Pharmacia], 0.25 unit/ml oligo(dT) [Pharmacia]; 0.05% Nonidet P40 and 10 mCi/ml 32 P-dTTP in the presence of various concentrations of test compound. After incubating for one hour at 37°C, 40 microliters of the reaction mixture was applied to a Schleicher & Schuell NA 45 membrane saturated with 2 x SSC (0.3 M NaCl, 30 mM sodium citrate, pH 7.0) in a Schleicher & Schuell Minifold over one sheet of GB003 filter paper. Each well of the minifold was washed four times with 2 x SSC. Autoradiography was performed, and radioactivity was quantified with a Packard Matrix (Meriden, CT) 9600 direct beta counter.

Cell Fusion Assay

20 Cell fusions were performed as described in Matthews *et al.*, *Proc. Natl. Acad. Sci. USA* 84:5424-5428 (1987). MOLT-4 cells (7×10^4) were incubated with HIV-1_{LAI} chronically infected CEM cells (10^4) in 96-well half-area flat-bottomed plates (Costar) in 100 microliters culture medium were incubated with the cell mixtures at 37°C for 24 hours. Multinucleated syncytia were enumerated by microscopic examination of the entire contents of each well.

25 The results of the above tests are shown in Table 1.

TABLE 1
Anti-HIV, HIV-RT, and Fusion Assay for
Betulinic Acid and Dihydrobetulinic Acid Derivatives

Anti-HIV Assay			HIV-RT Assay	Fusion Assay
Compounds	EC ₅₀ (μM) ^a	T.I.	IC ₅₀ (μg/mL) ^b	IC ₁₀₀ (μg/mL) ^c
1	1.4	9.3	>100	>100
2	2.7	5.9	>100	20 ^d
3	3.5 x 10 ⁻⁴ - 1 x 10 ⁻⁵	20,000 - 700,000	>100	20 ^d
4	2.3 x 10 ⁻³	1,974.0	>100	30 ^d
5	0.01	1,172.0	>100	40 ^e
6	0.9	14.0	>100	>100
7	0.56	13.8	N.T. ^g	N.T. ^g
8	3.5 x 10 ⁻⁴ - 5 x 10 ⁻⁵	14,000 - 98,000	>100	20 ^f
9	5.7 x 10 ⁻³	1,017.0	>100	20 ^f
10	5.6 x 10 ⁻³	2,344.0	>100	40 ^e
12	0.03 ^h	>90 ^h	No Inhibition ^h	3 ^h
a: Concentration which inhibits viral replication in H9 cells by 50% b: Concentration required to inhibit 50% of HIV-1 RT activity c: Concentration required to completely inhibit HIV- I induced syncytia mediated by CEM4 d: No noticeable cytotoxicity at 100 μg/mL e: Toxic to the cells at 80 μg/mL in one-day fusion assay f: Toxic at 40 μg/mL under the assay conditions g: N.T. = Not Tested h: Data from reference 8				

Additional tests were conducted on several compounds according to the present invention for inhibition of HIV-1 replication in H9 lymphocytes and PBMCs. These results are shown in Table 2.

TABLE 2
Inhibition of HIV-1 Replication in H9 Lymphocytes and PBMCs


Compound	Cell	EC ₅₀ (mg/mL)	TI ₅₀
8	H9	0.0032	17,188
8	PBMCs	0.00175	2,286
3	H9	0.00128	3,125
3	PBMCs	0.0175	229
10	H9	0.0032	2,344
5	H9	0.01	1,500

The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without departing from the generic concept, and therefore such adaptations and modifications are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation.

All references cited in this specification are hereby incorporated by reference.

What Is Claimed Is:




1. A compound of the formula:




wherein R is a C₂-C₂₀ mono- or di-carboxyacetyl group that may be optionally substituted, and

R' is one of hydrogen, C₂-C₁₀ substituted or unsubstituted alkyl, or aryl.

2. A compound according to claim 1, wherein R is selected from the group consisting of




or




-26-

3. A compound of the formula:



wherein R is a C₂-C₂₀ mono- or dicarboxyacyl groups that may be optionally substituted, and R' is hydrogen, C₂-C₁₀ substituted or unsubstituted alkyl, or aryl.

4. A compound according to claim 3, wherein R is selected from the group consisting of:



5. A pharmaceutical composition comprising a compound according to either claim 1 or claim 3, or a pharmaceutically acceptable ester, salt, ether, sulfate, or glucuronide thereof, and a pharmaceutically acceptable carrier.

6. A pharmaceutical composition according to claim 5, further comprising a drug selected from an anti-viral agent or an immunostimulating agent.

7. A pharmaceutical composition according to claim 6, wherein said antiviral agent is selected from the group consisting of gamma globulin, amantadine, guanidine, hydroxybenzimidazole, interferon- α , interferon- β , interferon- γ , thiosemicarbazones, methisazone, rifampin, ribavirin, pyrimidine analogs, purine analogs, foscarnet, phosphonoacetic acid, acyclovir, dideoxynucleosides, and gancyclovir.

8. A method for inhibiting a retroviral infection in cells or tissue of an animal comprising administering an effective retroviral inhibiting amount of a pharmaceutical composition according to claim 5.

9. The method according to claim 8, wherein said composition is administered to provide said compound in an amount ranging from about 0.1 to about 100 mg/kg body weight.

10. The method according to claim 9, wherein said composition is administered to provide said compound in an amount ranging from about 1 to about 100 mg/kg body weight.

11. The method according to claim 8, wherein said animal is a human.

12. A method for treating a patient suffering from a retroviral related pathology, comprising administering to said subject a retroviral inhibiting effective amount of a pharmaceutical composition according to claim 5.

13. A method according to claim 12 wherein said retroviral related pathology is an HIV infection.

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US96/09485
A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 560/116, 194; 514/510

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 560/116, 194; 514/510

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	JOURNAL OF NATURAL PRODUCTS, Volume 57, number 2, issued February 1994, T. FUJIOKA ET AL., "Anti-AIDS agents, 11. Betulinic acid and platanic acid as anti-HIV principles from syzigium claviflorum, and the anti-HIV activity of structurally related triterpenoids" pages 243-247, see the entire document.	1,3,5,8 ----- 2,4
X -- Y	GB 1415601 B4A (BIOREX LABORATORIES LIMITED), 26 November 1975, see the entire document.	3,5 ---- 4

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

16 SEPTEMBER 1996

Date of mailing of the international search report

02 OCT 1996

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Barbara Badio

Telephone No. (703) 305-1235

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	COLLECTION OF CZECHOSLOVAK CHEMICAL COMMUNICATIONS, Volume 37, number 2, issued 28 December 1970, J. KLINOT ET AL., "Triterpenes. XXI. 3,4-Seco derivatives of betulinic acid" pages 603-609, see the entire document.	1-4
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCE, volume 91 number 9, issued April 1994, MAYAUX ET AL., "Triterpene derivative that block entry of human immunodeficiency virus type 1 into cells" pages 3564-3568, see the entire document.	1, 5, 8


A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):


A01N 37/00; A61K 31/21; C07C 69/34, 69/74

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 451/02, A01N 43/40	A1	(11) International Publication Number: WO 96/37494 (43) International Publication Date: 28 November 1996 (28.11.96)
(21) International Application Number: PCT/GB96/01151 (22) International Filing Date: 13 May 1996 (13.05.96) (30) Priority Data: 9510459.2 24 May 1995 (24.05.95) GB (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): URCH, Christopher, John [GB/GB]; 40 Dalcross, Crown Wood, Bracknell, Berkshire RG12 3UJ (GB). SALMON, Roger [GB/GB]; 38 Tawfield, Bracknell, Berkshire RG12 4YU (GB). LEWIS, Terence [GB/GB]; 18 Quintilis, Roman Hill, Bracknell, Berkshire RG12 4QQ (GB). GODFREY, Christopher, Richard, Ayles [GB/GB]; 159 Viking, Great Hollands, Bracknell, Berkshire RG12 8UW (GB). CLOUGH, Martin, Stephen [GB/GB]; 25 Stephen Close, Twyford, Berkshire RG10 0XN (GB). (74) Agents: BISHOP, Nigel, Douglas et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealott's Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG42 6YA (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: BICYCLIC AMINES AS INSECTICIDES

(I)



(A)

(57) Abstract

A compound of formula (I), wherein R¹ represents a group of formula (A) where each of W, X, Y and Z represents either a group CR or the nitrogen atom, provided that not more than two of W, X, Y and Z represent the nitrogen atom and where each R present is independently selected from hydrogen and halogen atoms and cyano, amino, hydrazino, acylamino, hydroxy, alkyl, hydroxyalkyl, alkoxy, haloalkyl, haloalkoxy, alkenyl, alkenyloxy, alkoxyalkenyl, alkynyl, carboxylic acyl, alkoxycarbonyl, aryl and heterocyclyl groups, said groups comprising up to 6 carbon atoms, and wherein R² represents hydrogen or cyano or a group selected from alkyl, aryl, heteroaryl, aralkyl, heteroarylalkyl, alkenyl, aralkenyl, alkynyl, alkoxycarbonyl, alkanesulfonyl, arenesulfonyl, alkanyloxycarbonyl, aralkyloxycarbonyl, aryloxycarbonyl, heterocyclyalkyl, carbamyl or dithiocarboxyl groups, said groups comprising from 1 to 15 carbon atoms, said groups being optionally substituted with one or more substituents selected from, halogen, cyano carboxyl, carboxylic acyl, carbamyl, alkoxycarbonyl, alkoxy, alkylenedioxy, hydroxy, nitro, haloalkyl, alkyl, amino, acylamino, imidate and phosphonato groups; and acid addition salts and quaternary ammonium salts and N-oxide derived therefrom. The compounds are useful as insecticides.

oxygen or a group NR⁴), provided that when R is alkenyl, aralkenyl or alkynyl said group does not have an unsaturated carbon atom bonding directly to the ring nitrogen of formula (I); R³ and R⁴ are, independently, hydrogen, alkyl, aryl, heteroaryl, aralkyl, heteroarylalkyl, alkenyl, aralkenyl, alkynyl, heterocyclylalkyl, alkoxycarbonyl or carboxylic acyl; alkyl moieties of R, R³ and R⁴ comprise from 1 to 15 carbon atoms, and are optionally substituted with one or more substituents selected from halogen, cyano, carboxyl, carboxylic acyl, carbamyl, alkoxycarbonyl, alkoxy, alkylenedioxy, hydroxy, nitro, amino, acylamino, imidate and phosphonato groups; aryl, heteroaryl, aralkyl, heteroarylalkyl, alkenyl, aralkenyl, alkynyl, alkoxycarbonyl, alkanesulfonyl, arenesulfonyl, alkanyloxy carbonyl, aralkyloxy carbonyl, aryloxy carbonyl, heterocyclylalkyl, carbamyl or dithiocarboxyl moieties of R, R³ and R⁴ comprise from 1 to 15 carbon atoms, and are optionally substituted with one or more substituents selected from, halogen, cyano, carboxyl, carboxylic acyl, carbamyl, alkoxycarbonyl, alkoxy, alkylenedioxy, hydroxy, nitro, haloalkyl, alkyl, amino, acylamino, imidate and phosphonato groups; R¹ represents hydrogen, hydroxy, alkyl, alkoxy, amino, nitro, isocyanato, acylamino, hydroxyalkyl, optionally substituted heteroaryl, alkoxyalkyl, haloalkyl, haloalkoxyalkyl, aralkyloxyalkyl, acyloxyalkyl, amidoximido, sulfonyloxyalkyl, aminoalkyl, alkoxycarbonylamino, acylaminoalkyl, cyanoalkyl, imino, formyl, acyl or carboxylic acid or an ester or amide thereof, or alkenyl or alkynyl either of which is optionally substituted by halogen, alkoxy, cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl or cyano; or an acid addition salt, quaternary ammonium salt or N-oxide derived therefrom; an insecticidal, acaricidal or nematocidal composition comprising a compound of formula (I) and a suitable carrier or diluent therefore; a method of combating and controlling insect, acarine or nematode pests at a locus which comprises treating the pests or the locus of the pests with an effective amount of a compound of formula (I) or a composition as hereinbefore described.